













- Strathie LW, Wood AR, van Rooi C and McConnachie A (2005). *Parthenium hysterophorus* (Asteraceae) in southern Africa, and initiation of biological control against it in South Africa. In: Prasad TVR, Nanjappa HV, Devendra R, Manjunath A, Subramanya SC, Chandrashekar VK, Kiran Kumar KA, Jayaram KA and Prabhakara Setty TK (Eds) *Proc Second Int Conference on Parthenium Management*, pp 127-133. University of Agricultural Sciences, Bangalore, India.
- Terblanche C (2014a). National Strategy for the management of Famine Weed (*Parthenium hysterophorus*) in South Africa 2015-2018. Department of Environmental Affairs, Pretoria.
- Terblanche C (2014b). National Implementation Plan for the management of Famine Weed (*Parthenium hysterophorus*) in South Africa 2015-2016. Department of Environmental Affairs, Pretoria.
- Terblanche C, Nänni I, Kaplan H, Strathie LW, McConnachie AJ, Goodall J and van Wilgen BW (2016). An approach to the development of a national strategy for controlling invasive alien plant species: The case of *Parthenium hysterophorus* in South Africa. *Bothalia: African Biodiversity and Conservation*: in press.
- Tomley AJ (2000). *Puccinia melampodii* (summer rust), a new biocontrol agent for parthenium weed. In: Swarbrick JT (Ed), *Proc 6th Queensland Weed Symposium*, Queensland, Australia, pp. 126-129. Weed Science Society of Queensland, Brisbane.
- van der Laan M, Reinhardt CF, Belz RG, Truter WF, Foxcroft LC and Hurlle K (2008). Interference potential of the perennial grasses *Eragrostis curvula*, *Panicum maximum* and *Digitaria eriantha* with *Parthenium hysterophorus*. *Trop Grasslands* 42: 88-95.
- Wood AR and Scholler M (2002). *Puccinia abrupta* var. *partheniicola* on *Parthenium hysterophorus* in southern Africa. *Plant Dis* 86: 327.